



OPEN ACCESS

Top 10 research priorities for congenital diaphragmatic hernia in Australia: James Lind Alliance Priority Setting Partnership

Roberto Chiletto,^{1,2} Courtney Vodopic,³ Emiko Hunt,³ Jess Lawer,³ Monique Bertinetti,^{1,2} Stephanie Malarbi,^{1,4} Valerie Kyritsis,^{5,6} Scott Petersen,⁷ David Stewart,^{1,2} Jean Hellstern,² Michael Stewart ,^{1,4} Leah Hickey ,^{1,2} David G Tingay ,^{2,4} Trisha M Prentice ,^{1,2}

► Additional supplemental material is published online only. To view, please visit the journal online (<https://doi.org/10.1136/archdischild-2024-327108>).

¹The Royal Children's Hospital, Melbourne, Victoria, Australia

²Murdoch Children's Research Institute, Melbourne, Victoria, Australia

³Congenital Diaphragmatic Hernia Australia, Geelong, Victoria, Australia

⁴The University of Melbourne, Melbourne, Victoria, Australia

⁵La Trobe University, Melbourne, Victoria, Australia

⁶Monash University, Melbourne, Victoria, Australia

⁷Mater Mothers' Hospital, South Brisbane, Queensland, Australia

Correspondence to

Dr Trisha M Prentice;
trisha.prentice@mcri.edu.au

Received 5 March 2024

Accepted 19 April 2024

Published Online First

16 June 2024



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY. Published by BMJ Group.

To cite: Chiletto R, Vodopic C, Hunt E, et al. *Arch Dis Child Fetal Neonatal Ed* 2025;**110**:F23–F29.

ABSTRACT

Objectives The Gaps in the Congenital Diaphragmatic Hernia (CDH) Journey Priority Setting Partnership (PSP) was developed in collaboration with CDH Australia, James Lind Alliance (JLA) and the Murdoch Children's Research Institute to identify research priorities for people with CDH, their families and healthcare workers in Australasia.

Design Research PSP in accordance with the JLA standardised methodology.

Setting Australian community and institutions caring for patients with CDH and their families.

Patients CDH survivors, families of children born with CDH (including bereaved) and healthcare professionals (including critical care physicians and nurses (neonatal and paediatric), obstetric, surgical, allied health professionals (physiotherapists, speech pathologists and speech therapists) and general practitioners.

Main outcome measure Top 10 research priorities for CDH.

Results 377 questions, from a community-based online survey, were categorised and collated into 50 research questions. Through a further prioritisation process, 21 questions were then discussed at a prioritisation workshop where they were ranked by 21 participants (CDH survivors, parents of children born with CDH (bereaved and not) and 11 multidisciplinary healthcare professionals) into their top 10 research priorities.

Conclusion Stakeholders' involvement identified the top 10 CDH-related research questions, spanning from antenatal care to long-term functional outcomes, that should be prioritised for future research to maximise meaningful outcomes for people with CDH and their families.

INTRODUCTION

Congenital diaphragmatic hernia (CDH) is a life-threatening developmental defect of the diaphragm resulting in the herniation of abdominal contents into the thorax. This impacts lung and heart development, resulting in varying degrees of lung hypoplasia, pulmonary hypertension and/or left ventricle hypoplasia. Additionally, CDH is associated with other major anomalies in 30% of cases including cardiac and gastrointestinal.^{1,2}

The incidence of CDH in Australia is approximately 1:2500 pregnancies, with a reported

WHAT IS ALREADY KNOWN ON THIS TOPIC

⇒ Congenital diaphragmatic hernia (CDH) is a life-threatening condition requiring multicentre research collaboration to advance outcomes throughout the CDH journey.

WHAT THIS STUDY ADDS

⇒ We describe the top 10 CDH-related research questions as defined through an Australian Research Priority Setting Partnership involving healthcare professionals and the CDH community.

HOW THIS STUDY MIGHT AFFECT RESEARCH, PRACTICE OR POLICY

⇒ Research collaborations targeting the defined research priorities for CDH are likely to have the greatest impact for children born with CDH throughout their lifetime and their families.

survival rate for all cases (including antenatal deaths and terminations) averaging between 50% and 60%.³ Antenatal diagnosis, standardisation of perinatal and postnatal medical and surgical management, protective ventilation strategies, advances in pulmonary hypertension and cardiac therapies and extracorporeal life-support techniques are contributing factors for improved survival over the last two decades.^{4,5} However, children with CDH experience prolonged hospitalisation and long-term morbidities, impacting their quality of life into adulthood.⁶

Despite attempts to standardise CDH management, significant differences remain, internationally.⁷ Existing international consensus guidelines are mainly based on case reviews and consensus expert opinions, reflecting the challenges in creating high-grade knowledge within high-risk, high-acuity and relatively rare critical care populations.^{8,9} The primary focus of medical research in CDH has been on short-term markers of disease, with limited focus on long-term and functional outcomes,¹⁰ despite as many as 87% of CDH survivors experiencing long-lasting morbidity, including pulmonary, gastrointestinal and neurodevelopmental problems.^{11–14} The CDH journey, therefore, requires both interventions that optimise disease-free survival in the

Protected by copyright. Including for uses related to text and data mining, AI training, and similar technologies.

Arch Dis Child Fetal Neonatal Ed: first published as 10.1136/archdischild-2024-327108 on 16 June 2024. Downloaded from <http://fn.bmj.com/> on May 22, 2025 at Department GEZ-LTA ErasmusHogeschool.




Figure 1 Process diagram. PSP, priority setting partnership.

intensive care unit (ICU) and coordinated multidisciplinary and evidence-based care from the prenatal period through adulthood to reduce long-term morbidity.

Research in the setting of rare diseases requires costly and time-consuming multicentre collaboration. Therefore, research that delivers value to the end user should be prioritised. Research priority setting partnerships (PSPs) address these challenges by ensuring research questions with the potential to have the greatest impact for patients and families are identified. The ‘Gaps in the CDH (Australia) Journey PSP’ aimed to identify, prioritise and share the needs and questions of people with CDH, their families or carers and healthcare professionals to inspire future research, influence future care and improve outcomes and the well-being of those diagnosed with CDH.

METHODS

CDH Australia (CDHA) and the Murdoch Children’s Research Institute (MCRI) collaborated on the project, supported by the James Lind Alliance (JLA), a not-for-profit association developed to bring together patients, families and clinicians into PSP.¹⁵ CDHA, a not-for-profit organisation with more than 1000 members in Australia and New Zealand, assists and supports families affected by CDH; conducting and funding education and research within Australia. CDHA funded this project and MCRI provided in-kind support.

Standardised JLA methodology¹⁵ was used, which is summarised in figure 1.

PSP start-up (13 July 2022)

The steering group overseeing the PSP was established in July 2022 in accordance with the JLA guidelines. Its 14 members included CDHA representatives (each with lived and/or living experience of CDH) and multidisciplinary healthcare professionals (experts in the management of children affected by CDH). The CDHA Board invited families with lived or living experience to represent the CDH community. The steering group was led by a neonatologist (TMP) with expertise in managing newborns with CDH and qualitative research. A JLA adviser (Tamara Rader) supported the process in alignment with the values of the JLA, providing fair and transparent processes, and fostering respectful and balanced conversations. The MCRI assigned a project coordinator (JH) for organisational support. Meetings were held virtually to ensure broad representation.

PSP scope

The steering group was responsible for all aspects of the PSP design and scope. The scope included:

- Management of pregnancies where CDH has been antenatally detected.
- Counselling and communication to families about management of CDH.
- Postnatal management, including resuscitation, stabilisation, surgical intervention and postoperative management.
- Longer-term follow-up and management of patients with CDH; care of expectant parents previously diagnosed with CDH.
- Improving the well-being of patients and families who have been directly impacted by a diagnosis or care of a child with CDH.

The PSP limited the scope to services relevant to Australian practice.

Setting partnerships

Research stakeholders were identified by mapping out the CDH journey for children affected by CDH and their families, spanning the fetal–maternal unit, perinatal management, neonatal hospital care and long-term follow-up services. A diverse cultural and sociodemographic voice was prioritised and proactively sought, including review by RCH Wadja Aboriginal Family services. All steering group members contributed to the design and refinement of the first survey, including inviting members from their relevant communities to pilot the survey, with feedback incorporated prior to distribution.

Distribution of first survey (13 November 2022)

The first survey aimed to capture raw unanswered questions. Respondents could ask up to five open questions relevant to the CDH journey and were asked to provide demographic data, including connection to CDH (see online supplemental file A).

The survey was built on the REDCap¹⁶ web-based research data capture platform. The survey was launched on CDHA Sunflower Sunday (13 November 2022), an annual event hosted by CDHA and promoted with links via CDHA’s and MCRI’s websites and social media channels. It was also shared through medical institutions across Australia, Australian medical colleges and research institutions, appropriate email lists, social media channels and at national conferences. The survey closed on 31 March 2023.

Identifying unanswered questions

Questions submitted via the first survey were categorised into themes by members of the steering group. Small working groups of 2–3 steering group members were assigned a theme and collated questions into summary questions using non-technical language. Questions were excluded if answers existed within a Cochrane review or systematic review of randomised control trials (RCTs), or considered partially answered if a high-quality RCT or multicentre cohort study was completed. An electronic database search was performed of the Cochrane Database of Systematic Reviews, Ovid Medline and Embase using combinations and variations of predefined keywords. The results from the databases were merged and duplicates were removed. Only papers with full-text, published in English in the last 10 years were included for pragmatic review of whether sufficient uncertainty remained regarding the proposed research questions as per the JLA guidelines. Existing international guidelines were not considered sufficient to exclude a question. The remaining 50 summary questions were carried forward for the following prioritisation stage.

Distribution of the second survey (19 June 2023)

The second REDCap survey was advertised and distributed through the same channels as the first survey and directly to participants from the first survey who requested contact. Participants were asked to choose and rank their top 10 CDH-related research questions from the list of 50. Recipients also provided demographic data. Participants were randomised by computer allocation to receive one of three randomly ordered question lists to reduce bias from survey fatigue. The survey remained open for 1 month. Responses were listed and categorised in order, based on the number of overall preferences and relative ranking, with subanalysis reviewing differences between preferences for the lived experience community versus healthcare professionals. Following review by the steering group, the top 21 questions were brought forward to the final workshop for discussion.

Final workshop and top 10 questions

The workshop was held online and divided into two sessions on 2 consecutive days (2 and 3 August 2023). An expression of interest to participate in the workshop was distributed with additional selective targeting of under-represented stakeholder groups within the second survey. A total of 25 voting participants (and a support person) agreed to participate. Ultimately, two healthcare professionals (representing anaesthetic and obstetric maternal–fetal medicine) and three persons with lived experience withdrew due to personal reasons. Of the 21 who attended, 11 were healthcare professionals and 10 CDH community members (table 1). Each participant received, via post and email, a prereading package containing information on the workshop process and the list of 21 questions to be ranked in order of priority.

During the first workshop session, 4 evenly represented subgroups discussed the 21 questions and collaboratively proposed a new agreed-upon priority order, with assistance from an experienced JLA facilitator.

The priority order lists were integrated and an aggregated ranked list of questions was generated for the second session. During this session, four different subgroups were asked to discuss the aggregated list of questions from the plenary session to allow for new perspectives and final prioritisation. The final top 10 CDH research priorities in Australia were established

| Table 1 Workshop participants | | | |
|--|--|-----|----------|
| Member 1 | Raising a child with CDH | VIC | Regional |
| Member 2 | Bereaved mother | QLD | Regional |
| Member 3 | Bereaved mother | VIC | Regional |
| Member 4 | Raising a child with CDH | NSW | Metro |
| Member 5 | CDH survivor | NSW | Metro |
| Member 6* | Raising a child with CDH | WA | Metro |
| Member 7 | CDH survivor supported by mother (member 26) | QLD | Regional |
| Member 8 | Raising a child with CDH | VIC | Regional |
| Member 9* | Bereaved mother | WA | Metro |
| Member 10 | Bereaved mother | VIC | Metro |
| Member 11* | Raising two children with CDH | QLD | Regional |
| Member 12 | Raising a child with CDH | NSW | Metro |
| Member 13 | Clinical nurse consultant | QLD | Metro |
| Member 14 | Neonatologist | QLD | Metro |
| Member 15 | Neonatal clinical nurse | QLD | Metro |
| Member 16 | Physiotherapist | QLD | Metro |
| Member 17 | Transport NETS WA | WA | Regional |
| Member 18 | Paediatric surgeon | QLD | Metro |
| Member 19† | Paediatric anaesthetist | VIC | Metro |
| Member 20 | Senior paediatric dietician | VIC | Metro |
| Member 21 | PICU/ECMO | VIC | Metro |
| Member 22 | Respiratory paediatrician | VIC | Metro |
| Member 23 | Respiratory paediatrician | NSW | Metro |
| Member 24 | Midwife | NSW | Metro |
| Member 25‡ | Obstetric MFM | QLD | Regional |
| Member 26 | Clinical nurse and mother of CDH survivor (attended in support capacity to member 7) | QLD | Regional |
| *CDH-affected individuals/families who could not attend on the day. †Invited anaesthetic medical representatives who could not attend on the day. ‡Invited obstetric, maternal–fetal medicine who could not attend on the day. CDH, congenital diaphragmatic hernia; ECMO, extracorporeal membrane oxygenation; MFM, maternal–fetal medicine; NETS, newborn emergency transport service; NSW, New South Wales; PICU, paediatric intensive care unit; QLD, Queensland; VIC, Victoria; WA, Western Australia. | | | |

from the aggregated subgroup lists (see table 2). As per the JLA processes, participants were surveyed on their experience of participation.

RESULTS

First survey

377 questions were submitted by 175 respondents: 39.4% submitted 1 question, 20.6% 2, 12.6% 3, 6.3% 4 and 21.1% 5 questions.

Healthcare professionals and lived experience respondents comprised 55.5% and 44.5%, respectively, with neonatologists representing 43.8% of the medical professionals. The demographic characteristics are summarised in table 3, with a predominance of Caucasian women aged between 25 and 60 years living in a metropolitan area.

Interim prioritisation

From the original 377 questions 54 were excluded for being out of scope or unanswerable (eg, comments specific to an individual, or non-CDH specific). The remaining 317 questions were ordered into 8 categories: pathophysiology (16), antenatal (76), resuscitation (13), admission (75), follow-up (89), well-being

Table 2 Final top 10 questions and subgroup (G1–G4) ranking

| Rank | Question | G1 | G2 | G3 | G4 |
|------|--|----|----|----|----|
| 1 | How can we optimise the neurodevelopmental outcomes of survivors of CDH? | 1 | 1 | 1 | 1 |
| 2 | How does CDH impact feeding and gut health and how can these outcomes be improved? | 2 | 2 | 2 | 2 |
| 3 | What follow-up and surveillance should patients with CDH receive (through childhood and adulthood)? | 6 | 3 | 3 | 3 |
| 4 | How can we best support the immediate care (transition) of babies born with CDH? | 4 | 6 | 6 | 5 |
| 5 | Which infants with CDH would benefit from ECMO and what is the optimal timing? | 9 | 5 | 5 | 4 |
| 6 | What are the best strategies for managing pulmonary hypertension in CDH and when should they be used? | 3 | 8 | 7 | 7 |
| 7 | What are the predictors of long-term outcomes in patients with CDH? | 10 | 4 | 4 | 8 |
| 8 | How can we best support lung growth and function of patients with CDH during their initial hospital admission? | 7 | 7 | 9 | 6 |
| 9 | What (if any) antenatal interventions improve outcomes? | 5 | 10 | 8 | 10 |
| 10 | What are the long-term respiratory outcomes of CDH? | 11 | 12 | 11 | 11 |
| 11 | How does CDH affect the growth and development of other organs before and after birth? | 12 | 13 | 12 | 9 |
| 12 | How do we best support the mental health and well-being of parents and survivors of CDH? | 8 | 9 | 16 | 18 |
| 13 | How do the lungs of babies with CDH grow and develop with time? | 14 | 14 | 13 | 13 |
| 14 | Can stem cells improve the growth of the underdeveloped or affected lung in CDH? | 17 | 11 | 10 | 17 |
| 15 | What are the best antenatal predictors of outcome in CDH and how are they best measured (eg, MRI, ultrasound)? | 13 | 19 | 19 | 12 |
| 16 | What is the best way to support the breathing of babies born with CDH before surgical repair? | 16 | 17 | 15 | 16 |
| 17 | How should follow-up care be best coordinated for patients with CDH? | 15 | 15 | 14 | 21 |
| 18 | How should labour and delivery best be supported for babies with CDH? | 18 | 16 | 17 | 14 |
| 19 | What causes or worsens CDH antenatally? | 19 | 18 | 18 | 15 |
| 20 | How can we optimise sedation and pain relief throughout the patient's admission? | 20 | 20 | 20 | 19 |
| 21 | Does choice of inotropic medication to support blood pressure affect outcome for patients with CDH? | 21 | 21 | 21 | 20 |

G1–G4=subgroups 1–4.

Blue represents top 10 responses, yellow represents next 11–16 responses and red represents 17–21 responses.

CDH, congenital diaphragmatic hernia; ECMO, extracorporeal membrane oxygenation.

(14), next-generation (6) and implementation service provision (28).

Research questions with similar content were combined to form 65 summary questions. These were checked against 2446 published articles on CDH and 50 unanswered questions were submitted for prioritisation through the second survey.

Second survey

108 recipients completed the survey. Healthcare professionals represented 64.2% of the respondents, with 49.0% being neonatal physicians. As per the first survey, the predominant respondents were female, Caucasian and living in a metropolitan area. The top 21 ranked questions were taken to the final workshop.

Final workshop

The final workshop included 21 participants from 4 states/territories in Australia including regional areas: 10 CDH community members (1 in a support capacity) and 11 healthcare professionals (table 1).

Table 2 summarises the final top 10 research priorities in CDH as determined during the workshop, together with individual subgroup ranking (groups 1–4). 14 participants (14; 64% persons with lived experience, 36% healthcare professionals) responded to a postworkshop feedback survey; all either agreed or strongly agreed that the facilitators were fair and impartial, and 91% either agreed or strongly agreed the process was fair and robust.

DISCUSSION

CDH remains one of the most complex conditions managed in perinatal medicine, yet this is the first report of research priorities in CDH using a partnership approach. The results exemplify

the importance and validity of consumer participation in driving research. The CDHA community provided visibility to the experience of living with a condition with life-long implications to specialists who may deal with only an isolated aspect of CDH.

The majority of published research and consensus guidelines focus on critical care management of CDH, including optimal ventilatory settings and/or management of pulmonary hypertension.^{7 8 17} However, the top three questions prioritised by this PSP focused on the long-term quality of life of children with CDH. This highlights the CDH journey beyond the ICU with anticipated long-term multisystem morbidity including altered lung mechanics, scoliosis, gastro-oesophageal reflux¹⁸ and neurodevelopmental challenges such as risks of lower intellectual ability, motor challenges, autism spectrum disorder and attention deficit hyperactivity disorder.^{12 13} A recent review of the ‘unsolved problems in CDH follow-up’ summaries these into four main groups: identification of risk factors for longer-term morbidities, such as gastro-oesophageal reflux disease (GORD), correlation between prenatal predictors and late outcomes, neurodevelopmental and optimal surgical approaches based on patient characteristics.¹⁹ In contrast to this PSP, this ‘problem’ list was created by clinicians alone, yet encouragingly there is considerable overlap with the CDH PSP top 10 research priorities in the prioritisation of neurodevelopmental outcomes, longer-term morbidities and early prediction of these outcomes. Though the need for long-term multidisciplinary follow-up is recognised in international guidelines and consensus statements, evidence about the best composition, frequency and duration of follow-up is lacking.²⁰ Small case-loads at most institutions (15–20 babies per year) highlight the need to direct international multi-centred studies towards finding solutions to the issues of greatest priority, adjusting the scope from research on acute management to longer-term follow-up and morbidity.

Table 3 Demographic characteristic of the survey participants

| N (%) | 1st survey | 2nd survey |
|-----------------------------------|------------|------------|
| Valid responses | 175 | 108 |
| Respondent relation | | |
| CDH survivor | 5 (3.5) | 3 (3.2) |
| Parent/caregiver CDH survivor | 46 (31.9) | 23 (24.2) |
| Bereaved parent of CDH child | 9 (6.3) | 7 (7.4) |
| Healthcare professional | 80 (55.6) | 61 (64.2) |
| Other | 4 (2.8) | 1 (1.1) |
| Not disclosed | 31 | 13 |
| Healthcare provider role | | |
| NICU nurse | 14 (17.5) | 14 (23) |
| Midwife | 1 (1.3) | 0 |
| Nurse other specialty | 1 (1.3) | 0 |
| NICU doctor | 35 (43.8) | 30 (49.2) |
| Surgeon | 6 (7.5) | 1 (1.6) |
| Obstetrician | 8 (10) | 1 (1.6) |
| General paediatrician | 0 | 0 |
| General practitioner | 1 (1.3) | 0 |
| Doctor other specialty | 8 (10) | 6 (9.8) |
| Sonographer | 0 | 0 |
| Dietician | 1 (1.3) | 1 (1.6) |
| Speech pathologist | 2 (2.6) | 0 |
| Physiotherapist | 1 (1.3) | 0 |
| Occupational therapist | 1 (1.3) | 1 (1.6) |
| Social worker | 0 | 2 (3.3) |
| Psychologist | 1 (1.3) | 2 (3.3) |
| Other | 0 | 3 (4.9) |
| Gender | | |
| Male | 36 (25) | 27 (29) |
| Female | 103 (71.5) | 64 (68.8) |
| Non-binary | 1 (0.7) | 0 |
| Not disclosed | 4 (2.8) | 2 (2.2) |
| Age (years) | | |
| <15 | 0 | 0 |
| 15–24 | 3 (2.1) | 2 (2.1) |
| 25–40 | 60 (41.7) | 41 (43.2) |
| 41–60 | 71 (49.3) | 45 (47.4) |
| >60 | 6 (4.2) | 5 (5.3) |
| Not disclosed | 4 (2.8) | 2 (2.1) |
| Ethnicity | | |
| Aboriginal/Torres Strait Islander | 1 (0.7) | 1 (1.1) |
| African | 0 | 0 |
| Asian | 11 (7.6) | 6 (6.4) |
| Caucasian | 112 (77.8) | 83 (88.3) |
| Hispanic or Latino | 0 | 0 |
| Maori | 0 | 0 |
| Pacific Islander | 0 | 0 |
| Not disclosed | 12 (8.3) | 2 (2.1) |
| Other | 8 (5.6) | 2 (2.1) |
| Living area | | |
| Australian Capital Territory | 4 (2.8) | 3 (3.2) |
| New South Wales | 28 (19.4) | 12 (12.6) |
| Northern Territory | 0 | 0 |
| South Australia | 3 (2.1) | 2 (2.1) |
| Tasmania | 2 (1.4) | 0 |
| Queensland | 21 (14.6) | 15 (15.8) |
| Victoria | 57 (39.6) | 50 (52.6) |

Continued

Table 3 Continued

| N (%) | 1st survey | 2nd survey |
|-------------------|------------|------------|
| Western Australia | 14 (9.7) | 5 (5.3) |
| New Zealand | 10 (6.9) | 5 (5.3) |
| Other | 5 (3.5) | 3 (3.2) |
| Living setting | | |
| Metropolitan | 108 (75) | 72 (75.8) |
| Regional city | 30 (20.8) | 18 (19.8) |
| Remote | 6 (4.2) | 5 (5.3) |

CDH, congenital diaphragmatic hernia; NICU, neonatal intensive care unit.

Research PSPs bring together diverse stakeholders that allow for new perspectives and a shared commitment to considering what will have the greatest impact for those directly impacted by the condition. A clinician with experience in two PSPs in two different roles commented that as a patient, he ‘was attracted towards choosing support and rehabilitation priorities’ and as a healthcare professional, he ‘gravitated towards pathogenetic, diagnostic and treatment priorities’.²¹ In this PSP, for example, clinicians came to appreciate the significant negative impact of gastrointestinal complications and long-term feeding: all four subgroups ranked it independently as their number two priority, acknowledging that long-term incidence of gastro-oesophageal reflux for children with CDH remains high (up to 70%) and feeding difficulties persist beyond the first 5 years of life.^{6 22}

Another priority was evidence on the best short-term and long-term management approaches to pulmonary hypertension and, correlated to it, patient candidacy and timing for extra-corporeal membrane oxygenation (ECMO). Pulmonary hypertension is a well-recognised complication in CDH patients and is a major contributor to early mortality, impacts admission duration and is associated with adverse neurodevelopmental outcomes.^{23–25} Unlike ventilatory support, there have been few advances in effective treatments for pulmonary hypertension though most are crude and designed to manage acute pulmonary hypertensive crises. Future research is needed on the role of medium-term and long-term pulmonary hypertensive therapies, such as endothelin antagonists or prostacyclin, especially as point-of-care cardiac function monitoring has become standard practice.^{17 26} ECMO is rarely used in the management of infants born with CDH in Australia and is usually reserved as rescue therapy when pulmonary hypertensive, cardiac and high-frequency ventilation support have been maximised. ECMO timing and indication remain uncertain for clinicians and families, and variability still exists internationally on ECMO patient selection, timing of mechanical support and CDH repair while on or post-ECMO.²⁷

It was observed by the facilitators that some participants prioritised questions with the most immediate results and impact; research questions perceived to be ‘novel’ but ‘blue-sky thinking’ with slow anticipated translation into clinical practice (eg, stem cell therapies) were more likely to be prioritised lower. As JLA methodology excludes participants who do not identify as a lived experience expert or clinician, it is possible that the perspectives of translational CDH researchers are not reflected in the results of this PSP. Both evidence and research prioritisation may need regular review. Questions that were less specific to CDH (eg, parental mental health) were also prioritised lower, perhaps in the hope that evidence may be readily translatable from research in other conditions.

Limitations

Online and emailed surveys generally achieve lower response rates than face-to-face surveys and may present some initial selection bias, though diverse and representative participation was a core priority. Despite these efforts, we acknowledge that the predominance of Caucasian women participants, aged 25–60 years living in metropolitan areas, may have introduced some bias to the results.

The second survey had a high representation in the healthcare cohort from Victoria and NICU, which the steering group worked to balance in the final workshop with increased participation of interstate representatives and limited presence of neonatal staff (doctors or nurses). The high representations did not appear to affect priority congruence.

Under-representation from minority groups is a known challenge worldwide, due to mistrust, time and cost constraints, and cultural factors, such as language and communication.²⁸ Furthermore, unlike the UK and the USA, Australia lacks a robust system to collect race and ethnicity data to address under-representation and promote active participation at all levels of research funding.²⁹

Although this survey focused on the Australian experience, the journey of parents and children with CDH is likely similar across other specialty centres internationally. We therefore hypothesise that the priorities identified by this PSP are generally applicable to CDH populations in regions with similar healthcare models and systems.

CONCLUSION

Research PSP for CDH in Australia was a robust and well-accepted process by all stakeholders in determining the top 10 research priorities for CDH. Given the high congruence across healthcare professionals, and survivors and families with lived and living experience of CDH, directing research funding and resources to address these important questions spanning from antenatal care to long-term follow-up is likely to result in the greatest impact and improved outcomes for patients and their families.

X Leah Hickey @lmhickey

Acknowledgements The authors would like to thank Susan Fisher, Val Manuelpillai, Caitlin Thouliss and Nadine Griffiths for their contributions to the PSP steering group at various points throughout the partnership. Thank you to Tamara Rader (JLA facilitator) for her support of this project throughout.

Contributors CV, DT, EH, LH and TMP were instrumental in developing the concept and setting up the steering group. All steering group members contributed to the survey designs and refinement of the research questions. DS, LH, MS and TMP reviewed the literature for existing evidence. RC, TMP, CV, EH and LH wrote the first draft and all authors contributed to redrafting the manuscript. TMP is guarantor.

Funding This study is supported by CDH Australia and the Victorian Government Operational Infrastructure Support Program (Melbourne, Australia). DT is supported by a National Health and Medical Research Council Leadership Level 1 Investigator Grant (Grant ID 2008212). TMP is supported by the RCH Foundation Clinical Scientist Fellowship.

Competing interests None declared.

Patient consent for publication Not applicable.

Ethics approval This study involves human participants. Human research ethics committee (HREC) approval was waived following institutional review by the Royal Children's Hospital (RCH) given the focus on research planning and stakeholder engagement. The RCH deemed that HREC approval was not required as this is a stakeholder consultation process rather than research participation, exempted this study. Participants gave informed consent to participate in the study before taking part.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available up reasonable request.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Michael Stewart <http://orcid.org/0000-0003-2235-4814>

Leah Hickey <http://orcid.org/0000-0001-8732-1453>

David G Tingay <http://orcid.org/0000-0003-1522-4691>

Trisha M Prentice <http://orcid.org/0000-0003-4271-7950>

REFERENCES

- Edel GG, Schaaf G, Wijnen RMH, et al. Cellular origin(S) of congenital diaphragmatic hernia. *Front Pediatr* 2021;9:804496.
- Politis MD, Bermejo-Sánchez E, Canfield MA, et al. Prevalence and mortality in children with congenital diaphragmatic hernia: a multicountry study. *Ann Epidemiol* 2021;56:61–9.
- O'Mahony E, Stewart M, Sampson A, et al. Perinatal outcome of congenital diaphragmatic hernia in an Australian tertiary hospital. *Aust N Z J Obstet Gynaecol* 2012;52:189–94.
- Holden KI, Harting MT. Recent advances in the treatment of complex congenital diaphragmatic hernia—a narrative review. *Transl Pediatr* 2023;12:1403–15.
- Gupta VS, Harting MT, Lally PA, et al. Mortality in congenital diaphragmatic hernia: a multicenter Registry study of over 5000 patients over 25 years. *Ann Surg* 2023;277:520–7.
- Peetsold MG, Heij HA, Kneepkens CMF, et al. The long-term follow-up of patients with a congenital diaphragmatic hernia: a broad spectrum of morbidity. *Pediatr Surg Int* 2009;25:1–17.
- Jancelewicz T, Brindle ME, Guner YS, et al. Toward standardized management of congenital diaphragmatic hernia: an analysis of practice guidelines. *J Surg Res* 2019;243:229–35.
- Snoek KG, Reiss IKM, Greenough A, et al. Standardized postnatal management of infants with congenital diaphragmatic hernia in Europe: the CDH EURO consortium consensus - 2015 update. *Neonatology* 2016;110:66–74.
- Puligandla P, Skarsgard E, Baird R, et al. Diagnosis and management of congenital diaphragmatic hernia: a 2023 update from the Canadian congenital diaphragmatic hernia collaborative. *Arch Dis Child Fetal Neonatal Ed* 2024;109:239–52.
- Lewis L, Sinha I, Losty PD. Clinical trials and outcome reporting in congenital diaphragmatic hernia overlook long-term health and functional outcomes—A plea for core outcomes. *Acta Paediatr* 2022;111:1481–9.
- van den Hout L, Sluiter I, Gischler S, et al. Can we improve outcome of congenital diaphragmatic hernia? *Pediatr Surg Int* 2009;25:733–43.
- Gerall CD, Stewart LA, Price J, et al. Long-term outcomes of congenital diaphragmatic hernia: a single institution experience. *J Pediatr Surg* 2022;57:563–9.
- Danzer E, Hoffman C, D'Agostino JA, et al. Neurodevelopmental outcomes at 5 Years of age in congenital diaphragmatic hernia. *J Pediatr Surg* 2017;52:437–43.
- Gunn-Charlton JK, Burnett AC, Malarbi S, et al. Neonatal neuroimaging after repair of congenital diaphragmatic hernia and long-term neurodevelopmental outcome. *World Jnl Ped Surgery* 2020;2:e000037.
- Alliance JL. JLA guidebook version 10. 2021. Available: <https://www.jla.nihr.ac.uk/jla-guidebook/> [Accessed Dec 2023].
- Harris PA, Taylor R, Thielke R, et al. Research electronic data capture (REDCap)—A metadata-driven methodology and workflow process for providing translational research Informatics support. *J Biomed Inform* 2009;42:377–81.
- Yang MJ, Russell KW, Yoder BA, et al. Congenital diaphragmatic hernia: a narrative review of controversies in neonatal management. *Transl Pediatr* 2021;10:1432–47.
- Tan JK, Banton G, Minutillo C, et al. Long-term medical and psychosocial outcomes in congenital diaphragmatic hernia survivors. *Arch Dis Child* 2019;104:761–7.
- Valfré L, Conforti A, Morini F, et al. Unsolved problems in CDH follow-up. *Front Pediatr* 2022;10:977354.
- Puligandla PS, Skarsgard ED, Offringa M, et al. Diagnosis and management of congenital diaphragmatic hernia: a clinical practice guideline. *CMAJ* 2018;190:E103–12.
- Adebajo A. Patients and healthcare professionals in priority setting partnerships: two very different sides of the same coin. *BMJ* 2022;378:o2310.

- 22 Power B, Shibuya S, Lane B, *et al.* Long-term feeding issue and its impact on the daily life of congenital diaphragmatic hernia survivors: results of the first patient-led survey. *Pediatr Surg Int* 2020;36:63–8.
- 23 Montalva L, Raffler G, Riccio A, *et al.* Neurodevelopmental impairment in children with congenital diaphragmatic hernia: not an uncommon complication for survivors. *J Pediatr Surg* 2020;55:625–34.
- 24 Herco M, Sloan P, Vogel A, *et al.* Survival and neurodevelopmental outcomes in congenital diaphragmatic hernia patients with single versus repeat extracorporeal membrane oxygenation runs. *Am J Perinatol* 2022.
- 25 Grover TR, Rintoul NE, Hedrick HL. Extracorporeal membrane oxygenation in infants with congenital diaphragmatic hernia. *Semin Perinatol* 2018;42:96–103.
- 26 Ramaraj AB, Rice-Townsend SE, Foster CL, *et al.* Trends in use of prostacyclin analogs for management of CDH-associated pulmonary hypertension. *Pediatr Surg Int* 2022;38:1241–7.
- 27 Guner Y, Jancelewicz T, Di Nardo M, *et al.* Management of congenital diaphragmatic hernia treated with Extracorporeal life support: interim guidelines consensus statement from the Extracorporeal life support organization. *ASAIO J* 2021;67:113–20.
- 28 Shea L, Pesa J, Geonnotti G, *et al.* Improving diversity in study participation: patient perspectives on barriers, racial differences and the role of communities. *Health Expect* 2022;25:1979–87.
- 29 Renzaho AMN. The lack of race and Ethnicity data in Australia-A threat to achieving health equity. *Int J Environ Res Public Health* 2023;20:5530.

Gaps in the CDH Journey

Identifying research priorities to improve the outcomes of babies born with Congenital Diaphragmatic Hernia. The "Gaps in the CDH Journey" project is a collaboration between the Murdoch Children's Research Institute and CDH Australia, supported by the James Lind Alliance.

This project's vision is to improve the outcomes for children born with CDH through understanding the research priorities of people with a lived experience of Congenital Diaphragmatic Hernia in Australia. (What is CDH?)

The result will be the identification of the Top 10 Priorities for CDH Research in Australia.

The Top 10 Priorities will be published and shared with key groups such as research funders, healthcare providers, hospitals, research institutes, researchers, and the wider community.

By identifying, prioritising, and sharing the needs and questions of people with CDH, their families and healthcare providers our goal is to inspire further research, influence future care, and improve outcomes and wellbeing of those diagnosed with CDH in Australia.

This is your chance to have a direct and meaningful role in optimising the outcomes of babies diagnosed with, and those affected by CDH.

About this survey This online survey is the first stage of the "Gaps in the CDH Journey" project and will be used to collect CDH-related questions.

You can submit up to 5 questions relating to any aspect of the CDH journey, including diagnosis, antenatal preparation, postnatal treatment, and future implications.

All questions collected in this survey will be checked to determine if they have already been answered by existing research, which will also be shared at the conclusion of this project.

The questions not yet answered by existing research will form a second survey where those affected by CDH, caregivers, and healthcare professionals will choose and prioritise what they feel are the most important.

The outcome of this project will be the identification of the "Top 10 CDH Research Priorities".

The Top 10 Priorities will be published and shared with key groups such as research funders, healthcare providers, researchers, and the wider community.

By identifying, prioritising, and sharing the needs of those affected by CDH, our goal is to inspire further research, influence future care, and improve outcomes and wellbeing of those diagnosed with CDH in Australia.

It is anticipated this survey will take 10-15 minutes to complete.

Who can take part Families in the CDH community who have received medical care in Australia are invited to participate. Healthcare providers who work with babies and families or people affected by CDH in Australia are also encouraged to have their say.

We want to hear from you if your experience with CDH is based in Australia, and you are one of the following:

A CDH survivor A parent or caregiver to a child who has survived CDH A bereaved parent or caregiver to a child with CDH A healthcare professional who works with children, adults and/or families affected by CDH Will my answers be kept confidential? Yes, all submissions to this survey are confidential.

Your submission can be anonymous, meaning we do not know who you are, or you may choose to provide your contact details if you would like to participate in the second stage of the project or receive project updates.

By participating in this survey, you are agreeing to have your questions included in our "Gaps in the CDH Journey" project. Your questions will be analysed and may be published but will not be linked to you or any organisation or group you are affiliated with.

Any contact details collected as part of this survey will be stored confidentially and securely in accordance with the Privacy Act and deleted at the end of the project. MCRI Privacy Policy and The Privacy Act

More information Murdoch Children's Research Institute MCRI Research: Gaps in the CDH journey Email: jean.m.petersen@mcri.edu.au

projectredcap.org



CDH Australia Email: hello@cdh.org.au

James Lind Alliance

Your questions about CDH

You may submit up to 5 CDH-related questions that you would like future CDH research to address, or that highlight gaps in healthcare and management for CDH families.

Draw on your own experience and consider what is important to you.

Some areas you may want to consider are:

CDH diagnosis
Pregnancy care and preparation for the birth of a baby diagnosed with CDH
Birth and postnatal care for babies diagnosed with CDH
Future or ongoing physical and mental health and wellbeing implications for children born with CDH
Any other questions or areas for exploration

Remember these are just examples and should not limit you. Research can cover a wide variety of subjects, so anything you think of will be of interest. This is your chance to be heard and to inform research.

Select the number of questions you would like to submit?

☐ 1

☐ 2

☐ 3

☐ 4

☐ 5

(Please let us know what CDH-related questions are important to you by filling in the text boxes below. You may change the number of questions you submit by clicking on the numbers.)

My first question

My second question

My third question

My fourth question

My fifth question

About You

Please tell us a little about yourself. We will use this information to ensure we are reaching a wide range of people and to better understand the needs of different groups.

In relation to your CDH journey, which role best describes you?

- ☐ A CDH survivor
- ☐ Parent/caregiver of a CDH survivor
- ☐ Bereaved parent/caregiver of a child with CDH
- ☐ Healthcare professional
- ☐ Other

Please specify

What is your primary role as a Health Care Provider?

- ☐ NICU Nurse
- ☐ Midwife
- ☐ Nurse - other speciality
- ☐ NICU Doctor
- ☐ Surgeon
- ☐ Obstetrician
- ☐ General Paediatrician
- ☐ General Practitioner
- ☐ Doctor - other speciality
- ☐ Sonographer
- ☐ Dietician
- ☐ Speech Pathologist
- ☐ Physiotherapist
- ☐ Occupational Therapist
- ☐ Social Worker
- ☐ Psychologist
- ☐ Other

Please specify

Do you predominantly care for adult or paediatric patients with CDH?

- ☐ Children
- ☐ Adults

What is the gender you identify with?

- ☐ Female
- ☐ Male
- ☐ Non-binary
- ☐ Prefer not to say

What is your current age?

- ☐ < 15 years
- ☐ 15 - 24 years
- ☐ 25 - 40 years
- ☐ 41 - 60 years
- ☐ 61 + years
- ☐ Prefer not to say

What ethnicity do you identify with?

- ☐ Aboriginal/Torres Strait Islander
- ☐ African (including African American)
- ☐ Asian
- ☐ Caucasian
- ☐ Hispanic or Latino
- ☐ Maori
- ☐ Pacific Islander
- ☐ Prefer not to say
- ☐ Other

Please specify

Where do you currently live?

- ☐ ACT and other territories
- ☐ New South Wales
- ☐ Northern Territory
- ☐ South Australia
- ☐ Tasmania
- ☐ Queensland
- ☐ Victoria
- ☐ Western Australia
- ☐ New Zealand
- ☐ Other

Please specify

Which best describes where you live?

- ☐ Metropolitan
- ☐ Regional City
- ☐ Remote

Stay in touch

If you would like to participate in the second stage of the project, or receive project updates, please select “Yes” below and provide your contact details.

The second stage of the “Gaps in the CDH Journey” project will take place in 2023.

Participants in the second stage of the project, will help to prioritise the questions submitted in this survey to determine the Top 10 Priorities for CDH in Australia.

The results will be available at the end of 2023.

Would you like to be involved in the next survey to put the research questions in order of importance?

☐ Yes

☐ No

Would you like to be kept informed on the progress of the project and the results?

☐ Yes

☐ No

If you answered yes to either question, please let us know your contact details. Any contact details collected as part of this survey will be stored confidentially and securely in accordance with the Privacy Act and deleted at the end of the project.

Name

Email